




Blackleg in SA 2020

2020 SA Seasonal Conditions



Winter rainfall deciles (Jun-Aug)



Spring rainfall deciles (Sep-Nov)

2020 SA Blackleg Monitoring Site Data

SA	Group A, MR-MS	Group B, MR	Group C, MS	Group BF, MR-MS	Group AD, R-MR	Group ABDF, R	Group H, R
Arthurton	No recommendation as disease too low for both CSII and Leaf lesions						
Cummins	Yellow	Red	Red	Red	Green	Green	Green
Keith	Red	Red	Red	Red	Green	Green	
Riverton	Red	Red	Yellow	Red	Green	Green	
Roseworthy	Red	Yellow	Red	Red	Green	Green	
Spalding	No recommendation as disease too low for both CSII and Leaf lesions						
Wangary	Yellow	Red	Red	Green	Green	Green	
Yeelanna	Red	Red	Red	Red	Green	Green	Green

2019 Group H on Lower EP



Current Practice on Lower EP

- The majority of the LEP canola is limited to varieties with Group B resistance
 - but all MR+
- Some push away from Group 7 (SDHI) seed dressings
 - So Group 7 can be used as an foliar application 6-8 leaf
- A divide amongst commercial and independent agronomists on results from fungicide applications targeted at controlling UCI.